

Review

Unique roles in health promotion of dietary flavonoids through gut microbiota regulation: Current understanding and future perspectives



Zhanming Li ^{a,b,c}, Zhongyang Ren ^{b,*}, Lin Zhao ^c, Lin Chen ^d, Yue Yu ^a, Dongxu Wang ^a, Xuejin Mao ^e, Guangtian Cao ^f, Zhenlei Zhao ^g, Hongshun Yang ^{c,*}

^a School of Grain Science and Technology, Jiangsu University of Science and Technology, Zhenjiang 212100, Jiangsu, China

^b College of Ocean Food and Biological Engineering, Jimei University, Xiamen 361021, China

^c Department of Food Science and Technology, National University of Singapore, Singapore 117542, Singapore

^d Food, Nutrition and Health, Faculty of Land and Food Systems, The University of British Columbia, Vancouver, BC, Canada

^e State Key Laboratory of Food Science and Technology, Nanchang University, Nanchang 330047, China

^f College of Standardization, China Jiliang University, Hangzhou 310018, China

^g Zhejiang Provincial Key Lab of Geriatrics & Geriatrics Institute of Zhejiang Province, Department of Geriatrics, Zhejiang Hospital, Hangzhou 310013, China

ARTICLE INFO

Keywords:

Flavonoid

Dietary supplement

Gut microbiota

Bioaccessibility

Metabolism

ABSTRACT

Advances in understanding the biological effects of dietary flavonoids and flavonoid-rich foods have been reported. Improving knowledge about their beneficial effects, and mechanisms of action, is crucial for better utilization. However, mechanisms responsible for their health benefits are still unclear. Previous research considered has suggested that gut microbiota might be linked to the metabolism of dietary flavonoids. To understand the bioactivities of dietary flavonoids/flavonoid-rich foods better, and the role of microbiota, we explored systematically 1) types of dietary flavonoids and associated health benefits, 2) low bioaccessibilities and metabolic characteristics, 3) gut microbiota role in regulation, and 4) crosstalk between regulation mechanisms. Current challenges and future perspectives were also considered, offering new research directions and identifying trends in the development of flavonoid-rich food products.

1. Introduction

Flavonoids, as important phenolic compounds, are widely distributed in fruits and vegetables, and their health benefits have been studied intensively (Battino et al., 2021; Yang et al., 2021; Xiao, 2022). Flavonoids ingested with food enter the circulatory system through the intestinal tract to exert their beneficial effects (Wang, Zhao, Wen, Ho, & Li, 2021). Numerous studies suggest that dietary flavonoid-rich foods and flavonoid compounds present a diverse range of biological activities (Barreca et al., 2021; Zhang et al., 2022). However, the large-scale application of flavonoids as food dietary supplements are still limited due to their poor bioavailability, instability in gastrointestinal digestion, and lack of targeted bioaffinity (Teng, Zheng, Cao, Huang, Xiao, & Chen, 2021).

Generally, the substances absorbed are not in the original format but become metabolites (Joardder, Kumar, & Karim, 2018). Moreover, many studies did not focus on investigating their metabolites at a physiological concentration (Chen, Cao, Huang, Xiao, & Teng, 2021).

Recently, gut microbiota has been applied in the studies of the metabolism of dietary flavonoids (Duan et al., 2021; Pei, Liu, & Bolling, 2020). It has been demonstrated that dietary flavonoids and gut microbiota can together contribute to the production of catabolites and metabolites (Chen & Yang, 2020). The microbial populations could act as an indicator of the individual responses to food ingredients (Qu et al., 2018). The fermentation of different dietary flavonoids and other substances in the intestine produce different metabolites that affect the intestinal environment and regulate gut microbiota (Rowland et al., 2018; Yang, Zhu, Sun, Ai, Edirisuriya, & Zhou, 2020).

During the past decade, reports on flavonoid activities and regulatory mechanisms proliferated (Ferraz et al., 2020; Maaliki, Shaito, Pintus, El-Yazbi, & Eid, 2019; Maleki, Crespo, & Cabanillas, 2019; Yu, Pei, & Li, 2022). However, the mechanisms involved in the health benefits of dietary flavonoids are still unclear. Moreover, flavonoids and their metabolites may exert their unique activities through multiple biological pathways (Daveri et al., 2018). The investigation of crosstalk among the signal pathways is highly demanded to evaluate the bioactivities of

* Corresponding authors.

E-mail addresses: rzy0529@jmu.edu.cn (Z. Ren), fstyngs@nus.edu.sg (H. Yang).

dietary flavonoids.

In the current perspective, the main types of dietary flavonoids, their bioaccessibility properties, and the health benefits of dietary flavonoids or flavonoid-rich foods are overviewed. More importantly, the effects of dietary flavonoids on regulating gut microbiota are summarized to highlight their roles in health promotion. In addition, the challenges and further perspectives for clarifying the health benefits of dietary flavonoids are highlighted, and new research opportunities for precision nutrition based on dietary flavonoids are provided.

2. Health benefits of dietary flavonoids

2.1. Flavonoid bioaccessibility

Flavonoids are divided into different classes based on heterocycle types, and shown in Fig. S1 are the six major classes, namely flavanols, flavanones, flavones, anthocyanins, flavonols, and isoflavones (Dias, Pinto, & Silva, 2021). As the main phenolic compounds in nature, flavonoids are the most studied phenolic compounds due to their potential benefits (Williamson, Kay, & Crozier, 2018). The exertion of dietary flavonoids' health benefits is confronted with obstacles, especially the low bioaccessibility, which is attributed to the compounds' low stability and non-targeted release (Toro-Uribe et al., 2019). Therefore, improving the bioaccessibility of dietary flavonoids is crucial in promoting their health benefits (Ravisankar, Agah, Kim, Talcott, Wu, & Awika, 2019; Song et al., 2022).

Recent studies on the performance of flavonoid bioaccessibility *in vitro* digestion are summarized in Table 1. In fact, flavonoid bioaccessibility is associated with many factors, among which food matrix effects are important for the bioactivities of dietary flavonoids (Alashi, Taiwo, Oyedele, Adebooye, & Aluko, 2019). The absorption and excretion of flavonoids and metabolites are both associated with their ability to bind with macronutrients in foods (Kamiloglu, Tomas, Ozdal, & Capanoglu, 2021). The non-covalent interactions of flavonoids and proteins, carbohydrates, or lipids have attracted intensive attention. For instance, the interaction with macronutrients via van der Waals interactions could affect the flavonoids' properties, including bioaccessibility and radical scavenging activity (Meng et al., 2021). Generally, proteins, carbohydrates, lipids, and other components in the food matrix are all influential in dietary flavonoid digestion (Kamiloglu, Tomas, Ozdal, & Capanoglu, 2021).

To overcome poor bioaccessibility, delivery systems have been developed for the targeted release of flavonoids to retain their bioactivities, thereby solving the problem of low oral bioavailability and poor intestinal absorption for some flavonoids (Zhang, Zhu, Yang, Huang, & Ho, 2021). Liposomal encapsulation, and micro- and nano-encapsulation studies have been performed to determine the beneficial effects of flavonoids (Zhao et al., 2018). Nano-encapsulation is a promising technique to deliver various hydrophobic flavonoids *in vivo* (Ayala-Fuentes & Chavez-Santoscoy, 2021). Importantly, novel scientific reports are crucial to support future opportunities to reveal food-body interactions and to develop delivery systems with edible materials for dietary flavonoids, as well as other functional food products (McClements, 2020).

2.2. Dietary flavonoid metabolism

Flavonoids, flavonoid-rich foods, and their derivatives, should present their bioactivities as human dietary supplements to exert their health benefits (Wang, Zhao, Wen, Ho, & Li, 2021). Foods containing flavonoids can reach the gut lumen and then enter the bloodstream following further absorption and metabolism (Ribas-Agustí, Martín-Belloso, Soliva-Fortuny, & Elez-Martínez, 2018). For example, the catabolism of anthocyanins and flavanones produces many derivative compounds, including 4-protocatechuic acid, gallic acid, and vanillic acid (Kay, Pereira-Caro, Ludwig, Clifford, & Crozier, 2017). The

Table 1

In vitro changes in the bioaccessibility of flavonoids.

Flavonoid source	Processing	Bioaccessibility performance	Reference
Rutin in Meghalayan cherry pomace	Microwave-assisted extraction and <i>In vitro</i> SGD	More than 50 %	(Kashyap, Riar, & Jindal, 2022)
Flavonoids from <i>Diospyros kaki</i>	<i>In vitro</i> SGD	100 % higher than in aqueous leaf extract	(Martínez-Las Heras, Pinazo, Heredia, & Andrés, 2017)
Galloylated flavonoids from grape seeds extract	Sodium caseinate binding interaction	Increase in both foam and emulsion	(Elegbede, Li, Jones, Campanella, & Ferruzzi, 2018)
Naringenin from tomato sauce	Industrial processing	Overall positive effect	(Tomas, Beekwilder, Hall, Sagdic, Boyacioglu, & Capanoglu, 2017)
Proanthocyanidin in baobab fruit shell	<i>In vitro</i> SGD	Increase	(Ismail et al., 2021)
Flavonoids from six quinoa seeds	<i>In vitro</i> SGD	12.64 %	(Pellegrini et al., 2017)
Anthocyanins from <i>Carissa macrocarpa</i>	Freeze drying and <i>in vitro</i> SGD	Increase	(Seke, Manhivi, Shoko, Slabbert, Sultanbawa, & Sivakumar, 2021)
Yoghurt fortified with rutin-casein coprecipitate	<i>In vitro</i> SGD	Increase	(Acevedo-Fani, Ochoa-Grimaldo, Loveday, & Singh, 2021)
Flavonoids-enriched juice mixture	Processing of pulsed electric technologies	Increase	(Buniowska, Carbonell-Capella, Frigola, & Esteve, 2017)
Fruit juices	<i>In vitro</i> SGD	24.25 %–67.00 %	(Mihaylova, Desseva, Stoyanova, Petkova, Terzyiska, & Lante, 2021)
Flavonoids in finger millet	Roast with lime juice added	25 % increase	(Hithamani & Srinivasan, 2017)

Note: SGD-simulated gastrointestinal digestion.

absorption of flavonoids requires the removal of glycosides via hydrolysis. In addition, cytosolic β -glucosidase and lactase phlorizin hydrolase, are employed for glucoside hydrolyzation (Kulkarni et al., 2017; Yang, Liu, Yang, Gupta, & Jiang, 2018). After hydrolysis, many transporters and metabolizing enzymes contribute to flavonoid metabolism. Taking (–)-epicatechin metabolism as an example, the ingested (–)-epicatechin-3'-O- β -D-glucuronide is hydrolyzed for further conversion (Fig. 1). Different metabolites with varying abundances, including structurally related metabolites, 5-carbon rings, and carbon-side chain ring fission metabolites, were shown at different times (0–36 h) in systemic circulation and urine (Ottaviani et al., 2016).

Besides, dietary flavonoids can interact with glucuronic acid, methyl, and sulfate in intestinal phase II metabolism, and the related metabolites interfere with many biochemical processes (Chalet, Hollebrands, Janssen, Augustijns, & Duchateau, 2018). Fifteen metabolites of bioactive flavonoids from *Scutellaria baicalensis* Georgi (Lamiaceae) have been identified as sulfate/glucuronide conjugates, acetylated, deoxygenated, hydroxylated, and methylated products (Du et al., 2015). Moreover, the major metabolites of quercetin are phloroglucinol, dihydrocafeic acid, 2-(3,4-dihydroxyphenyl)acetic acid, and 3,4-dihydroxybenzoic acid (Di Pede et al., 2020; Tang et al., 2016). Importantly, low-molecular-weight phenolics can be transformed from flavonoids by colonic microbiota, and the absorption sites and forms of flavonoids are key factors in determining their metabolism and bioactivities (Catalkaya et al., 2020). Gallic acid, 3,4-dihydroxybenzoic acid, and 3-hydroxybenzoic acid




Fig. 1. (*-*)-Epicatechin metabolites with the most abundance in human after dietary intake (0–36 h). SREM: structurally related (*-*)-epicatechin metabolites. 5C-RFM: 5-C ring fission metabolites. 3/1C-RFM: 3-C-side and 1-C-side chain ring fission metabolites. Adapted from reference (Ottaviani et al., 2016) with permission.

derived from theaflavin-3-ogallate catabolism can be determined by microbiota-mediated steps as shown in Fig. S2 (Pereira-Caro et al., 2017). Moreover, the phase II metabolism and colonic microbiota metabolism of dietary flavonoids are different in the catabolism of dietary flavonoids (Baky, Elshahed, Wessjohann, & Farag, 2021; Williamson et al., 2018).

Moreover, the bioactivities of flavonoids may synergize or weaken the bioactivities of their parent flavonoids. The combination of natural extracts and apigenin-quercetin synergistically can inactivate ATP

binding cassette transporter expression and inhibits P-glycoprotein activity for enhancing their bioavailability (Ravisankar et al., 2019). Recently, a systematical review has been highlighted to summarize some potential pathways for the metabolism of representative flavonoids from a historical perspective, as an attractive work to enrich our knowledge of flavonoid metabolism (Williamson et al., 2018).




Fig. 2. Scheme for the metabolism of ingested flavonoid-rich foods. All the parameters were cited from (Kalt, McDonald, Liu, & Fillmore, 2017; Murota et al., 2018; Pei et al., 2020; Pereira-Caro et al., 2017; Williamson et al., 2018).

2.3. Health benefits of dietary flavonoids

Many factors influence the ability of dietary flavonoids to exert health benefits, including age, sex, food matrix, digestive enzymes, and intestinal transporters (Fig. 2) (Murota, Nakamura, & Uehara, 2018; Murphy, Walker, Dyer, & Bryan, 2019). Notably, the investigations of the metabolite profiles in different sites are remarkable to elucidate the mechanisms on the health promotion of dietary flavonoids and their derivatives. The appearance of radiolabeled compound profiles in urine, plasma, and other organs can provide information relevant to dietary flavonoid metabolism, their absorption sites, and forms. Fundamental data can be acquired using the isotopic label to describe the absorption and distribution of dietary flavonoids, as well as metabolism and excretion. The species-dependent differences in metabolites are also valuable for revealing the mechanisms of the health benefits, as well as strengthening the health-promotion investigations and safety assessments of dietary flavonoid intake (Nie et al., 2021; Ottaviani et al., 2016; Zeng et al., 2019).

Dietary flavonoid-rich foods are associated with the improvement of health status (Brewis, SturtzSreetharan, & Wutich, 2018; Wu & Ballantyne, 2020; Zhou et al., 2022). The health benefits of dietary flavonoids are involved in the amelioration of bone mineral density (Sarmadi, Ismail, Yusof, & Yunoh, 2020), acute lung injury (Tang et al., 2018), UV-induced skin damage (Liu et al., 2018), and depression (Khan, Perviz, Sureda, Nabavi, & Tejada, 2018).

Moreover, dietary flavonoid consumption produces substantial

health status improvement in risk populations (Rees, Dodd, & Spencer, 2018). Long-term dietary flavonoid intake is considerably involved in lower odds of subjective cognitive decline, as health-promotion effects, reported in a 16-year investigation in the US with the participation of 49,493 women and 27,842 men (Yeh, Yuan, Ascherio, Rosner, Willett, & Blacker, 2021). However, the conclusive direct evidence regarding flavonoid-induced health benefits is also limited. Clarifying the potential health benefits of long-term dietary flavonoid intake is still clearly urgent.

2.4. Crosstalk of signaling pathway regulation

The primary actions of flavonoids are due to their redox and physical binding activities. Flavonoids have been employed to regulate the NFE2-related factor 2/heme-oxygenase-1 (Nrf2/HO-1) pathway of HepG2 Cells *in vitro* and inhibit reactive oxygen species production (Yu, Li, Cao, Huang, & Yang, 2019). The effects of flavonoids on inflammatory response in cells are mediated by regulating the mitogen-activated kinase/nuclear factor kappa B (MAPK/NF- κ B) pathway in cells. Consequently, dietary flavonoids are used to potentiate the inhibition of IFN- α on cancer cell viability by activating the JAK2/STAT3 signaling pathways (Li et al., 2016). In addition, flavonoids have elevated the P38/PPAR- α pathway to inhibit cell apoptosis and autophagy in mice (Lu et al., 2018).

As shown in Fig. 3A, dietary flavonoids modulate other crosstalk points associated with NF- κ B and Nrf2 pathways (Williamson et al.,




Fig. 3. (A) Regulation of chemokines, enzymes, and crosstalk proteins by flavonoids via Nrf2 and NF- κ B pathways. Reprinted from the reference (Williamson et al., 2018) with permission. (B) Dietary flavonoid fermentation produces short-chain fatty acids to regulate lipid and glucose metabolism, immune responses, and mediate related gene expression. Reprinted from the reference (Gentile and Weir, 2018) with permission.

2018). NF-κB, Nrf2, MAPK, P13K, and Akt-mTOR signaling pathways, as well as multiple signaling pathways, have been revealed as important candidates for the modulation of flavonoid bioactivity (Chu et al., 2019; Jian et al., 2020; Kim, Cheong, Lee, Sok, & Kim, 2020; Yuan, Che, Zhang, & Ruan, 2021). The evidence of crosstalk between NF-κB and Nrf2 pathways comprises the association of Rac Family Small GTPase 1 (RAC1) with the Kelch-like ECH associated protein 1 (KEAP1)-NF-κB and Nrf2 pathways, attenuating inflammation and oxidative stress (Blaser, Dostert, Mak, & Brenner, 2016; Taniguchi & Karin, 2018). Certainly, flavonoid-rich foods and their metabolites may likely share common crosstalk among protein/enzymes and receptors to regulate multiple transcription factors and signaling pathways for exerting their mechanism of action (Ciumărnean et al., 2020; Williamson et al., 2018).

3. Roles of dietary flavonoids in gut microbiota regulation

Recently, the relationship between gut microbiota and health promotion has been a research focus (Soderborg et al., 2018; Wan et al., 2019). It has been considered that gut microbiota are involved in health status modulation (Forbes-Hernandez et al., 2016; Agus, Planchais, & Sokol, 2018; Fan & Pedersen, 2021). In this section, the roles of gut microbiota regulation by dietary flavonoid intake are highlighted to evaluate the health benefits, as well as short-chain fatty acids (SCFAs) regulation and bile acid metabolism. In addition, the crosstalk of signaling pathways is summarized to exhibit the regulation mechanism of dietary flavonoids for health promotion.

3.1. Gut microbiota reconstruction

The regulation of dietary flavonoids reshaping gut microbiota for health benefits is the frontier of food nutrition research (Annunziata et al., 2021; Zmora, Suez, & Elinav, 2019). Gut microbiota regulation mediated by dietary flavonoid intake is associated with health benefits, contributing to the regulation of host immunity, inflammatory status, and the alteration of microbiota diversity (Li, Christman, Li, & Gu, 2020). During the past decades, studies focused on increasing beneficial bacteria, inhibiting deleterious species, and exploring interactions between dietary flavonoids and gut microbiota (Ma et al., 2019).

The majority of gut microbiota associated with flavonoid

transformation contribute to the O-deglycosylation of flavonoids in flavonoid transformation, such as *Bifidobacteriaceae*, *Lactobacillaceae*, *Lachnospiraceae*, and *Enterococcaceae* (Kim et al., 2020; Yan & Zhang, 2022; Barber, Houghton, & Williamson, 2021), whereas others are involved in the cleaving of flavonoid-C-glucosides or the degradation of aglycones (Braune & Blaut, 2016).

Flavonoids can improve health status by reconstructing the gut microbiota. Bilberry anthocyanin extract intake promotes the intestinal barrier function by reshaping the gut microbiota (Li, Wu, Li, Wang, Chen, & Lyu, 2019). The fermentation in the intestine of different dietary structures produces different metabolites that affect the intestinal environment and regulate gut microbiota (Makki, Deehan, Walter, & Bäckhed, 2018). Diets with high flavonoid contents might contribute to the alteration of gut microbiota composition and function within weeks (Maier et al., 2017). Table 2 shows recent studies on the roles of flavonoids in gut microbiota regulation. In general, flavonoids are subject to catabolism by gut microbiota, and the mechanism is likely via the regulation of gut microbiota, although the regulatory mechanisms of certain flavonoids and extracts require further exploration.

The different roles of gut microbiota have been explored recently, and the host circadian rhythm can be regulated through the gut-brain axis and the improvement of gut microecology, supporting new clues to clarify the health benefits of dietary flavonoids (Guo et al., 2019). However, the gut microbiota-dietary flavonoid interactions associated with the circadian rhythm remain complex and unclear (Song, Yang, Zhang, & Wang, 2021). The framework is proposed to provide new insights to discover the roles of dietary flavonoids and gut microbiota (Fig. S3) for circadian rhythm regulation. More efforts should be performed to reveal the complex relationships between circadian rhythm, gut microbiota, and dietary flavonoids, especially from the perspective of the gut-liver axis and gut-brain axis.

3.2. Short-chain fatty acids and bile acid metabolism

Flavonoids usually play a similar role as gut prebiotics in regulating the metabolism of SCFAs (Qi et al., 2021). SCFAs are the metabolites of carbohydrates fermented by gut microbiota, such as acetic acid, propionic acid, butyric acid, isobutyric acid, valeric acid, and isovaleric acid (Sanna et al., 2019). Diet structures have reshaped the gut microbiota to

Table 2
Performance of dietary flavonoids in gut microbiota regulation.

Flavonoid	Model	Mechanism	Reference
Aglycone quercetin	Germ-free mice with non-alcoholic fatty liver	Decrease of <i>Akkermansia</i> abundance and increase of NLRP3 expression	(Porras et al., 2019)
Anthocyanin	Colorectal cancer animal model	Substantial reduction of pro-inflammatory <i>Bilophila wadsworthia</i>	(Fernández, García, Monte, Villar, & Lombó, 2018)
Isoorientin	BALB/c mice	Inhibited growth of inflammation-related bacteria	(Yuan, Li, He, Gao, Wang, & Shao, 2018)
Myricetin	Nonalcoholic fatty liver disease in rats	Modulations in fecal butyric-acid-related bacteria and improvement of gut barrier function	(Sun et al., 2021)
Baicalein	Rats with high-fat diet-induced diabetes	Increase of SCFA-producing bacteria and decrease of pathogens, Positive effects produced by <i>Bacteroidales S24-7</i>	(Zhang et al., 2018)
Flavonoids from whole-grain oat	High fat fed C57BL/6N mice	Decrease of <i>Colidextrabacter</i> , <i>Desulfovibrio</i> , and <i>Lachnoclostridium</i>	(Duan et al., 2021)
Grape seed proanthocyanidins	Female Wistar rats	Increase of <i>Bacteroidetes</i> and <i>Proteobacteria</i> , reduction of <i>Firmicutes</i>	(Casanova-Martí et al., 2018)
Flavonoids from <i>Cyclocarya paliurus</i>	High-fat diet-induced obesity mouse model	Decrease in the ratio of <i>Firmicutes/Bacteroidetes</i>	(Cheng et al., 2019)
Ogan juice	C57BL/6J mice with high-fat diet	Decrease of phylum <i>Erysipelatoclostridiaceae</i> and genus <i>Erysipelatoclostridium</i> , and remarkable increase of SCFA-producing bacteria	(Guo et al., 2021)
Flavonoid-rich extract from <i>Grifola frondosa</i>	Rats with high-fat diet	Increased abundance of <i>Butyrimonas</i> genus	(Pan et al., 2018)
Naringin	ApoE ^{-/-} mice with high-fat diet	Regulation of bile salt hydrolase- and 7α-dehydroxylase-producing bacteria levels	(Wang et al., 2020)
Cranberry extract	High fat/high sucrose-fed C57BL/6J mice	Increase of the population of <i>Akkermansia</i>	(Anhê et al., 2015)
Mulberry leaves	Obese mice with high-fat diet	Increase of <i>Akkermansia</i> level and <i>Bacteroidetes/Firmicutes</i> ratio	(Sheng et al., 2019)
Flavonoid-rich extract from Green Seaweed	Mice with Type 2 diabetes	Higher abundance of <i>Lachnospiraceae</i> and <i>Alistipes</i>	(Yan et al., 2019)

change the composition of SCFAs (Zhu, Han, Ding, Zhu, Song, & Xiao, 2021). Moreover, a high-fat diet causes a decrease in SCFAs compared with a low-fat diet (Wan et al., 2019). SCFAs are associated with many biochemical processes, playing an important role in health benefits. SCFAs have been used as an energy source for intestinal epithelial cells and signaling molecules for the peripheral organs of the intestine. They are also employed to promote the stable state of the gut microbiota (Gentile & Weir, 2018). Fig. 3B shows that SCFAs can regulate lipid metabolism, glucose metabolism, and immune responses, and mediate related gene expression.

Enterohepatic circulation is involved in the metabolism of dietary flavonoids. As an immune organ comprising immune cells, the liver is exposed to gut microbiota through the portal vein. The intimate cross-talk between the liver and the gut is associated with the metabolism of nutrients and the clearance of bacterial metabolites (Gentile and Weir, 2018; Koudoufio, Desjardins, Feldman, Spahis, Delvin, & Levy, 2020; Moorthy, Chaiyakunapruk, Jacob, & Palanisamy, 2020). Bile acid metabolism is a complex process, and the metabolites of microbial regulation contribute to the diversity of the hydrophobic bile acid pool. The regulation of gut microbiota might be associated with both bile acid synthesis and uptake. Moreover, dietary flavonoids can be used to promote bile acid homeostasis (Duan et al., 2021; Wang, Xia, Yang, & Tang, 2019).

4. Current challenges and future perspectives

Although much progress has been achieved to strengthen and clarify the metabolism, bioavailability, and health promotion of dietary flavonoids, it is difficult to verify the full potential of dietary flavonoids due to the numerous challenges and barriers to clarifying their roles. As the most susceptible property, the low bioavailability of dietary flavonoids contributes to the limitations of health benefits. The smart, flexible design of delivery systems or encapsulation with edible materials is urgently needed to address the low stability and non-targeted release of

dietary flavonoids.

Lifespan-essential dietary flavonoids are consistently associated with health promotion, and the lack of flavonoids leads to an increased risk of developing chronic diseases. Current studies are mainly performed using animal models or *in vitro* experiments, and medium- and long-term health promotion investigations should be strengthened for understanding the mechanisms of health regulation of dietary flavonoids or flavonoid-rich food intake.

As one of the frontiers of food nutrition, the reshaping of gut microbiota by dietary flavonoids for health promotion has attracted numerous interests. Engagingly, future food design should be performed according to gut microbiota feeding for achieving health benefits (Ercolini & Fogliano, 2018). Although much progress has been made on gut microbiota regulation by dietary flavonoids, achieving the goal of health benefits and precision nutrition is still an urgent task. Challenges remain for the identification of gut microbiota in complex matrices, particularly at low concentrations with insufficient resolution, as well as uncultured microbes. Although considerable efforts are underway currently by multi-omic analysis, advanced techniques should be further developed to evaluate the regulatory mechanisms of dietary flavonoids for gut microbiota.

Although numerous studies have indicated that different dietary flavonoids or metabolites and catabolites of flavonoids achieve health promotion through gut microbiota regulation, systematic research is still urgent to clarify the mechanism of action. Therefore, in the current perspective, a research framework (Fig. 4) is proposed, contributing to further research and understanding of the benefits of dietary flavonoids through gut microbiota regulation. It may also help perform innovative novel research toward precision nutrition.

Given the insufficient reports on the change of gut microbiota by dietary flavonoid intake among populations in different areas, achieving a distinct conclusion on the differences in gut microbes that contribute to flavonoid metabolism among populations in different areas is difficult. Gut microbiota among populations in different areas may vary due




Fig. 4. Research framework for the analysis of the health promotion mechanisms of dietary flavonoids through gut microbiota regulation, as well as the systematic research toward precision nutrition.

to their diet structures, causing dissimilar behaviors of flavonoid metabolism (Kawabata, Yoshioka, & Terao, 2019). Precision nutrition among populations in different areas may be developed by the clarification of gut microbiota that contributes to flavonoid metabolism (O'Grady & Shanahan, 2021).

Importantly, more valuable results can be achieved by performing research to (i) clarify the molecular mechanism of dietary flavonoids for gut microbiota regulation accurately, (ii) evaluate the metabolic changes of digestion under the modulation of food matrix–flavonoid interactions by using simulated digestion system, (iii) understand the mechanisms of dietary flavonoids under metabolic environments comprehensively rather than simply reflecting the output of health effects in human beings, (iv) reveal the theoretical molecular mechanisms of dietary flavonoids for gut microbiota regulation by multi-omic analysis and intelligent algorithms, (v) determine the individual response to dietary flavonoids in different intestinal microecological backgrounds, and (vi) develop precision nutrition based on dietary flavonoids-gut microbiota interactions.

Overall, the current perspectives provide new insights into how dietary flavonoids regulate gut microbiota and how gut microbiota improves health status, offering new prospects for future research trends and opportunities to boost functional food product development toward precision nutrition.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

No data was used for the research described in the article.

Acknowledgments

Financial support from the National Natural Science Foundation of China (No. 31902204) and Natural Science Foundation of Fujian Province (2021J01835) were appreciated.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.foodchem.2022.133959>.

References

- Acevedo-Fani, A., Ochoa-Grimaldo, A., Loveday, S. M., & Singh, H. (2021). Digestive dynamics of yoghurt structure impacting the release and bioaccessibility of the flavonoid rutin. *Food Hydrocolloids*, 111, Article 106215.
- Agus, A., Planchais, J., & Sokol, H. (2018). Gut microbiota regulation of tryptophan metabolism in health and disease. *Cell Host & Microbe*, 23(6), 716–724.
- Alashi, A. M., Taiwo, K. A., Oyedele, D. J., Adebooye, O. C., & Aluko, R. E. (2019). Polyphenol composition and antioxidant properties of vegetable oil-fortified bread. *Journal of Food Biochemistry*, 43(6), e12625.
- Anhé, F. F., Roy, D., Pilon, G., Dudonné, S., Matamoros, S., Varin, T. V., et al. (2015). A polyphenol-rich cranberry extract protects from diet-induced obesity, insulin resistance and intestinal inflammation in association with increased Akkermansia spp. population in the gut microbiota of mice. *Gut*, 64(6), 872–883.
- Annunziata, G., Sureda, A., Orhan, I. E., Battino, M., Arnone, A., Jiménez-García, M., et al. (2021). The neuroprotective effects of polyphenols, their role in innate immunity and the interplay with the microbiota. *Neuroscience & Biobehavioral Reviews*, 128, 437–453.
- Ayala-Fuentes, J. C., & Chavez-Santoscoy, R. A. (2021). Nanotechnology as a key to enhance the benefits and improve the bioavailability of flavonoids in the food industry. *Foods*, 10(11), 2701.
- Baky, M. H., Elshahed, M. S., Wessjohann, L. A., & Farag, M. A. (2021). Interactions between dietary flavonoids and the gut microbiome: A comprehensive review. *British Journal of Nutrition*, 1–46.
- Barber, E., Houghton, M. J., & Williamson, G. (2021). Flavonoids as human intestinal α-glucosidase inhibitors. *Foods*, 10(8), Article 1939.
- Barreca, D., Trombetta, D., Smeriglio, A., Mandarli, G., Romeo, O., Felice, M. R., et al. (2021). Food flavonols: Nutraceuticals with complex health benefits and functionalities. *Trends in Food Science & Technology*, 117, 194–204.
- Battino, M., Giampieri, F., Cianciosi, D., Ansary, J., Chen, X., Zhang, D., et al. (2021). The roles of strawberry and honey phytochemicals on human health: A possible clue on the molecular mechanisms involved in the prevention of oxidative stress and inflammation. *Phytomedicine*, 86, Article 153170.
- Blaser, H., Dostert, C., Mak, T. W., & Brenner, D. (2016). TNF and ROS crosstalk in inflammation. *Trends in Cell Biology*, 26(4), 249–261.
- Braune, A., & Blaut, M. (2016). Bacterial species involved in the conversion of dietary flavonoids in the human gut. *Gut Microbes*, 7(3), 216–234.
- Brewis, A., SturtzSreetharan, C., & Wutich, A. (2018). Obesity stigma as a globalizing health challenge. *Globalization and Health*, 14(1), 1–6.
- Buniowska, M., Carbonell-Capella, J. M., Frigola, A., & Esteve, M. J. (2017). Bioaccessibility of bioactive compounds after non-thermal processing of an exotic fruit juice blend sweetened with Stevia rebaudiana. *Food Chemistry*, 221, 1834–1842.
- Casanova-Martí, A., Serrano, J., Portune, K. J., Sanz, Y., Blay, M. T., Terra, X., et al. (2018). Grape seed proanthocyanidins influence gut microbiota and enteroendocrine secretions in female rats. *Food & Function*, 9(3), 1672–1682.
- Cataliaga, G., Venema, K., Lucini, L., Rocchetti, G., Delmas, D., Daglia, M., et al. (2020). Interaction of dietary polyphenols and gut microbiota: Microbial metabolism of polyphenols, influence on the gut microbiota, and implications on host health. *Food Frontiers*, 1(2), 109–133.
- Chalet, C., Hollebrands, B., Janssen, H.-G., Augustijns, P., & Duchateau, G. (2018). Identification of phase-II metabolites of flavonoids by liquid chromatography-ion-mobility spectrometry–mass spectrometry. *Analytical and Bioanalytical Chemistry*, 410(2), 471–482.
- Chen, L., Cao, H., Huang, Q., Xiao, J., & Teng, H. (2021). Absorption, metabolism and bioavailability of flavonoids: A review. *Critical Reviews in Food Science and Nutrition*, 1–13.
- Chen, T., & Yang, C. S. (2020). Biological fates of tea polyphenols and their interactions with microbiota in the gastrointestinal tract: Implications on health effects. *Critical Reviews in Food Science and Nutrition*, 60(16), 2691–2709.
- Cheng, L., Chen, Y., Zhang, X., Zheng, X., Cao, J., Wu, Z., et al. (2019). A metagenomic analysis of the modulatory effect of *Cyclocarya paliurus* flavonoids on the intestinal microbiome in a high-fat diet-induced obesity mouse model. *Journal of the Science of Food and Agriculture*, 99(8), 3967–3975.
- Chu, Q., Yu, X., Jia, R., Wang, Y., Zhang, Y., Zhang, S., et al. (2019). Flavonoids from *Apium americanum* mediкус leaves protect RAW264.7 cells against inflammation via inhibition of MAPKs, Akt-mTOR pathways, and Nrf2 activation. *Oxidative Medicine and Cellular Longevity*, 1563024.
- Ciumărnean, L., Milaciu, M. V., Runcan, O., Vesa, ř. C., Răchișan, A. L., Negrean, V., et al. (2020). The effects of flavonoids in cardiovascular diseases. *Molecules*, 25(18), 4320.
- Daveri, E., Cremonini, E., Mastaloudis, A., Hester, S. N., Wood, S. M., Waterhouse, A. L., et al. (2018). Cyanidin and delphinidin modulate inflammation and altered redox signaling improving insulin resistance in high fat-fed mice. *Redox Biology*, 18, 16–24.
- Di Pede, G., Bresciani, L., Calani, L., Petrangolini, G., Riva, A., Allegrini, P., et al. (2020). The human microbial metabolism of quercetin in different formulations: An in vitro evaluation. *Foods*, 9(8), 1121.
- Dias, M. C., Pinto, D. C., & Silva, A. M. (2021). Plant flavonoids: Chemical characteristics and biological activity. *Molecules*, 26(17), 5377.
- Du, L., Qian, D., Shang, E., Liu, P., Jiang, S., Guo, J., et al. (2015). UPLC-Q-TOF/MS-based screening and identification of the main flavonoids and their metabolites in rat bile, urine and feces after oral administration of *Scutellaria baicalensis* extract. *Journal of Ethnopharmacology*, 169, 156–162.
- Duan, R., Guan, X., Huang, K., Zhang, Y., Li, S., Xia, J., et al. (2021). Flavonoids from Whole-Grain oat alleviated high-fat diet-induced hyperlipidemia via regulating bile acid metabolism and gut microbiota in mice. *Journal of Agricultural and Food Chemistry*, 69(27), 7629–7640.
- Elegbede, J. L., Li, M., Jones, O. G., Campanella, O. H., & Ferruzzi, M. G. (2018). Interactions between flavonoid-rich extracts and sodium caseinate modulate protein functionality and flavonoid bioaccessibility in model food systems. *Journal of Food Science*, 83(5), 1229–1236.
- Ercolini, D., & Fogliano, V. (2018). Food design to feed the human gut microbiota. *Journal of Agricultural and Food Chemistry*, 66(15), 3754–3758.
- Fan, Y., & Pedersen, O. (2021). Gut microbiota in human metabolic health and disease. *Nature Reviews Microbiology*, 19(1), 55–71.
- Fernández, J., García, L., Monte, J., Villar, C. J., & Lombó, F. (2018). Functional anthocyanin-rich sausages diminish colorectal cancer in an animal model and reduce pro-inflammatory bacteria in the intestinal microbiota. *Genes*, 9(3), 133.
- Ferraz, C. R., Carvalho, T. T., Manchope, M. F., Artero, N. A., Rasquel-Oliveira, F. S., Fattori, V., et al. (2020). Therapeutic potential of flavonoids in pain and inflammation: Mechanisms of action, pre-clinical and clinical data, and pharmaceutical development. *Molecules*, 25(3), 762.
- Forbes-Hernandez, T. Y., Gasparrini, M., Afrin, S., Bompadre, S., Mezzetti, B., Quiles, J. L., et al. (2016). The healthy effects of strawberry polyphenols: Which strategy behind antioxidant capacity? *Critical Reviews in Food Science and Nutrition*, 56(sup1), 46–59.
- Gentile, C. L., & Weir, T. L. (2018). The gut microbiota at the intersection of diet and human health. *Science*, 362(6416), 776–780.
- Guo, X., Cao, X., Fang, X., Guo, A., & Li, E. (2021). Inhibitory effects of fermented Ougan (*Citrus reticulata* cv. *Suavissima*) juice on high-fat diet-induced obesity associated with white adipose tissue browning and gut microbiota modulation in mice. *Food & Function*, 12(19), 9300–9314.

- Guo, T., Ho, C. T., Zhang, X., Cao, J., Wang, H., Shao, X., et al. (2019). Oolong tea polyphenols ameliorate circadian rhythm of intestinal microbiome and liver clock genes in mouse model. *Journal of Agricultural and Food Chemistry*, 67(43), 11969–11976.
- Hithamani, G., & Srinivasan, K. (2017). Bioaccessibility of polyphenols from selected cereal grains and legumes as influenced by food acidulants. *Journal of the Science of Food and Agriculture*, 97(2), 621–628.
- Ismail, B. B., Guo, M., Pu, Y., Çavuş, O., Ayub, K. A., Watharkar, R. B., et al. (2021). Investigating the effect of in vitro gastrointestinal digestion on the stability, bioaccessibility, and biological activities of baobab (*Adansonia digitata*) fruit polyphenolics. *LWT*, 145, Article 111348.
- Jian, T., Chen, J., Ding, X., Lv, H., Li, J., Wu, Y., et al. (2020). Flavonoids isolated from loquat (*Eriobotrya japonica*) leaves inhibit oxidative stress and inflammation induced by cigarette smoke in COPD mice: The role of TRPV1 signaling pathways. *Food & Function*, 11(4), 3516–3526.
- Joarder, M. U., Kumar, C., & Karim, M. A. (2018). Prediction of porosity of food materials during drying: Current challenges and directions. *Critical Reviews in Food Science and Nutrition*, 58(17), 2896–2907.
- Kalt, W., McDonald, J. E., Liu, Y., & Fillmore, S. A. (2017). Flavonoid metabolites in human urine during blueberry anthocyanin intake. *Journal of Agricultural and Food Chemistry*, 65(8), 1582–1591.
- Kamiloglu, S., Tomas, M., Ozdal, T., & Capanoglu, E. (2021). Effect of food matrix on the content and bioavailability of flavonoids. *Trends in Food Science & Technology*, 117, 15–33.
- Kashyap, P., Riari, C. S., & Jindal, N. (2022). Effect of extraction methods and simulated in vitro gastrointestinal digestion on phenolic compound profile, bio-accessibility, and antioxidant activity of Meghalayan cherry (*Prunus nepalensis*) pomace extracts. *LWT*, 153, Article 112570.
- Kawabata, K., Yoshioka, Y., & Terao, J. (2019). Role of intestinal microbiota in the bioavailability and physiological functions of dietary polyphenols. *Molecules*, 24(2), 370.
- Kay, C. D., Pereira-Caro, G., Ludwig, I. A., Clifford, M. N., & Crozier, A. (2017). Anthocyanins and flavanones are more bioavailable than previously perceived: A review of recent evidence. *Annual Review of Food Science and Technology*, 8, 155–180.
- Khan, H., Perviz, S., Sureda, A., Nabavi, S. M., & Tejada, S. (2018). Current standing of plant derived flavonoids as an antidepressant. *Food and Chemical Toxicology*, 119, 176–188.
- Kim, N. Y., Cheong, S. H., Lee, K. J., Sok, D. E., & Kim, M. R. (2020). Anti-inflammatory effects of *Ribes diacanthum* Pall mediated via regulation of Nrf2/HO-1 and NF-κB signaling pathways in LPS-stimulated RAW 264.7 macrophages and a TPA-induced dermatitis animal model. *Antioxidants*, 9(7), 622.
- Koudoufio, M., Desjardins, Y., Feldman, F., Spahis, S., Delvin, E., & Levy, E. (2020). Insight into polyphenol and gut microbiota crosstalk: Are their metabolites the key to understand protective effects against metabolic disorders? *Antioxidants*, 9(10), 982.
- Kulkarni, T. S., Khan, S., Villagomez, R., Mahmood, T., Lindahl, S., Logan, D. T., et al. (2017). Crystal structure of β-glucosidase 1A from *T hermotoga neapolitana* and comparison of active site mutants for hydrolysis of flavonoid glucosides. *Proteins: Structure, Function, and Bioinformatics*, 85(5), 872–884.
- Li, H., Christman, L. M., Li, R., & Gu, L. (2020). Synergic interactions between polyphenols and gut microbiota in mitigating inflammatory bowel diseases. *Food & Function*, 11(6), 4878–4891.
- Li, J., Wu, T., Li, N., Wang, X., Chen, G., & Lyu, X. (2019). Bilberry anthocyanin extract promotes intestinal barrier function and inhibits digestive enzyme activity by regulating the gut microbiota in aging rats. *Food & Function*, 10(1), 333–343.
- Li, S., Yang, L., Wang, P., He, Y., Huang, J., Liu, H., et al. (2016). Dietary apigenin potentiates the inhibitory effect of interferon-α on cancer cell viability through inhibition of 26S proteasome-mediated interferon receptor degradation. *Food & Nutrition Research*, 60(1), 31288.
- Liu, Z., Hu, Y., Li, X., Mei, Z., Wu, S., He, Y., et al. (2018). Nanoencapsulation of cyanidin-3-O-glucoside enhances protection against UVB-induced epidermal damage through regulation of p53-mediated apoptosis in mice. *Journal of Agricultural and Food Chemistry*, 66(21), 5359–5367.
- Lu, X., Liu, T., Chen, K., Xia, Y., Dai, W., Xu, S., et al. (2018). Isorhamnetin: A hepatoprotective flavonoid inhibits apoptosis and autophagy via P38/PPAR-α pathway in mice. *Biomedicine & Pharmacotherapy*, 103, 800–811.
- Ma, H., Zhang, B., Hu, Y., Wang, J., Liu, J., Qin, R., et al. (2019). Correlation analysis of intestinal redox state with the gut microbiota reveals the positive intervention of tea polyphenols on hyperlipidemia in high fat diet fed mice. *Journal of Agricultural and Food Chemistry*, 67(26), 7325–7335.
- Maaliki, D., Shaito, A. A., Pintus, G., El-Yazbi, A., & Eid, A. H. (2019). Flavonoids in hypertension: A brief review of the underlying mechanisms. *Current Opinion in Pharmacology*, 45, 57–65.
- Maier, T. V., Lucio, M., Lee, L. H., VerBerkmoes, N. C., Brislawn, C. J., Bernhardt, J., et al. (2017). Impact of dietary resistant starch on the human gut microbiome, metaproteome, and metabolome. *MBio*, 8(5), e01343–e1317.
- Makkai, K., Deehan, E. C., Walter, J., & Bäckhed, F. (2018). The impact of dietary fiber on gut microbiota in host health and disease. *Cell Host & Microbe*, 23(6), 705–715.
- Maleki, S. J., Crespo, J. F., & Cabanillas, B. (2019). Anti-inflammatory effects of flavonoids. *Food Chemistry*, 299, Article 125124.
- Martínez-Las Heras, R., Pinazo, A., Heredia, A., & Andrés, A. (2017). Evaluation studies of persimmon plant (*Diospyros kaki*) for physiological benefits and bioaccessibility of antioxidants by in vitro simulated gastrointestinal digestion. *Food Chemistry*, 214, 478–485.
- McClements, D. J. (2020). Advances in nanoparticle and microparticle delivery systems for increasing the dispersibility, stability, and bioactivity of phytochemicals. *Biotechnology Advances*, 38, Article 107287.
- Meng, Y., Hao, L., Tan, Y., Yang, Y., Liu, L., Li, C., et al. (2021). Noncovalent interaction of cyanidin-3-O-glucoside with whey protein isolate and β-lactoglobulin: Focus on fluorescence quenching and antioxidant properties. *LWT*, 137, Article 110386.
- Mihaylova, D., Desseva, I., Stoyanova, M., Petkova, N., Terzyiska, M., & Lante, A. (2021). Impact of in vitro gastrointestinal digestion on the bioaccessibility of phytochemical compounds from eight fruit juices. *Molecules*, 26(4), 1187.
- Moorthy, M., Chaiyakunapruk, N., Jacob, S. A., & Palanisamy, U. D. (2020). Prebiotic potential of polyphenols, its effect on gut microbiota and anthropometric/clinical markers: A systematic review of randomised controlled trials. *Trends in Food Science & Technology*, 99, 634–649.
- Murota, K., Nakamura, Y., & Uehara, M. (2018). Flavonoid metabolism: The interaction of metabolites and gut microbiota. *Bioscience, Biotechnology, and Biochemistry*, 82(4), 600–610.
- Murphy, K. J., Walker, K. M., Dyer, K. A., & Bryan, J. (2019). Estimation of daily intake of flavonoids and major food sources in middle-aged Australian men and women. *Nutrition Research*, 61, 64–81.
- Nie, H., Chen, H., Li, G., Su, K., Song, M., Duan, Z., et al. (2021). Comparison of flavonoids and phenylpropanoids compounds in Chinese water chestnut processed with different methods. *Food Chemistry*, 335, Article 127662.
- O'Grady, J., & Shanahan, F. (2021). Macronutrients, microbiome and precision nutrition. *Current Opinion in Gastroenterology*, 37(2), 145–151.
- Ottaviani, J. I., Borges, G., Momma, T. Y., Spencer, J. P., Keen, C. L., Crozier, A., et al. (2016). The metabolome of [2-14 C](-)-epicatechin in humans: Implications for the assessment of efficacy, safety and mechanisms of action of polyphenolic bioactives. *Scientific Reports*, 6(1), 1–10.
- Pan, Y. Y., Zeng, F., Guo, W. L., Li, T. T., Jia, R. B., Huang, Z. R., et al. (2018). Effect of Grifola frondosa 95% ethanol extract on lipid metabolism and gut microbiota composition in high-fat diet-fed rats. *Food & Function*, 9(12), 6268–6278.
- Pei, R., Liu, X., & Bolling, B. (2020). Flavonoids and gut health. *Current Opinion in Biotechnology*, 61, 153–159.
- Pellegrini, M., Lucas-Gonzalez, R., Fernández-López, J., Ricci, A., Pérez-Álvarez, J. A., Sterzo, C. L., et al. (2017). Bioaccessibility of polyphenolic compounds of six quinoa seeds during in vitro gastrointestinal digestion. *Journal of Functional Foods*, 38, 77–88.
- Pereira-Caro, G., Moreno-Rojas, J. M., Brindani, N., Del Rio, D., Lean, M. E., Hara, Y., et al. (2017). Bioavailability of black tea theaflavins: Absorption, metabolism, and colonic catabolism. *Journal of Agricultural and Food Chemistry*, 65(26), 5365–5374.
- Porras, D., Nistal, E., Martínez-Flórez, S., Olcoz, J. L., Jover, R., Jorquera, F., et al. (2019). Functional interactions between gut microbiota transplantation, quercetin, and high-fat diet determine non-alcoholic fatty liver disease development in germ-free mice. *Molecular Nutrition & Food Research*, 63(8), 1800930.
- Qi, Z., Yi, L., Li, J., Cui, Y., Rehman, F. U., Yang, J., et al. (2021). The romance between gut microbiota and flavonoids extract from *Smilax Glabra* roxb. and its potent alleviation of fatty liver. *Food & Function*, 12, 7836–7850.
- Qu, L., Ren, J., Huang, L., Pang, B., Liu, X., Liu, X., et al. (2018). Antidiabetic effects of *Lactobacillus casei* fermented yogurt through reshaping gut microbiota structure in type 2 diabetic rats. *Journal of Agricultural and Food Chemistry*, 66(48), 12696–12705.
- Ravisankar, S., Agah, S., Kim, H., Talcott, S., Wu, C., & Awika, J. (2019). Combined cereal and pulse flavonoids show enhanced bioavailability by downregulating phase II metabolism and ABC membrane transporter function in Caco-2 model. *Food Chemistry*, 279, 88–97.
- Rees, A., Dodd, G. F., & Spencer, J. P. (2018). The effects of flavonoids on cardiovascular health: A review of human intervention trials and implications for cerebrovascular function. *Nutrients*, 10(12), 1852.
- Ribas-Agustí, A., Martín-Belloso, O., Soliva-Fortuny, R., & Elez-Martínez, P. (2018). Food processing strategies to enhance phenolic compounds bioaccessibility and bioavailability in plant-based foods. *Critical Reviews in Food Science and Nutrition*, 58 (15), 2531–2548.
- Rowland, I., Gibson, G., Heinken, A., Scott, K., Swann, J., Thiele, I., et al. (2018). Gut microbiota functions: Metabolism of nutrients and other food components. *European Journal of Nutrition*, 57(1), 1–24.
- Sanna, S., van Zuydam, N. R., Mahajan, A., Kurilshikov, A., Vila, A. V., Vösa, U., et al. (2019). Causal relationships among the gut microbiome, short-chain fatty acids and metabolic diseases. *Nature Genetics*, 51(4), 600–605.
- Sarmadi, B., Ismail, A., Yusof, L., & Yunoh, M. F. M. (2020). Mechanism of action of cocoa on bone metabolism in calcium-and estrogen-deficient rat model of osteoporosis: Evidence for site and dose-related responses and involvement of IGF-I. *Journal of Functional Foods*, 66, Article 103793.
- Seke, F., Manhivi, V. E., Shoko, T., Slabbert, R. M., Sultanbawa, Y., & Sivakumar, D. (2021). Effect of freeze drying and simulated gastrointestinal digestion on phenolic metabolites and antioxidant property of the natal plum (*Carissa macrocarpa*). *Foods*, 10(6), 1420.
- Sheng, Y., Liu, J., Zheng, S., Liang, F., Luo, Y., Huang, K., et al. (2019). Mulberry leaves ameliorate obesity through enhancing brown adipose tissue activity and modulating gut microbiota. *Food & Function*, 10(8), 4771–4781.
- Soderborg, T. K., Clark, S. E., Mulligan, C. E., Janssen, R. C., Babcock, L., Ir, D., et al. (2018). The gut microbiota in infants of obese mothers increases inflammation and susceptibility to NAFLD. *Nature Communications*, 9(1), 1–12.
- Song, D., Yang, C. S., Zhang, X., & Wang, Y. (2021). The relationship between host circadian rhythms and intestinal microbiota: A new cue to improve health by tea polyphenols. *Critical Reviews in Food Science and Nutrition*, 61(1), 139–148.

- Song, J., Yu, Y., Chen, M., Ren, Z., Chen, L., Fu, C., et al. (2022). Advancement of protein- and polysaccharide-based biopolymers for anthocyanin encapsulation. *Frontiers in Nutrition*, 9, Article 938829.
- Sun, W. L., Li, X. Y., Dou, H. Y., Wang, X. D., Li, J. D., Shen, L., et al. (2021). Myricetin supplementation decreases hepatic lipid synthesis and inflammation by modulating gut microbiota. *Cell Reports*, 36(9), Article 109641.
- Tang, Y., Nakashima, S., Saiki, S., Myoi, Y., Abe, N., Kuwazuru, S., et al. (2016). 3, 4-Dihydroxyphenylacetic acid is a predominant biologically-active catabolite of quercetin glycosides. *Food Research International*, 89, 716–723.
- Tang, X., Zhao, H., Jiang, W., Zhang, S., Guo, S., Gao, X., et al. (2018). Pharmacokinetics and pharmacodynamics of citrus peel extract in lipopolysaccharide-induced acute lung injury combined with Pinelliae Rhizoma Praeparatum. *Food & Function*, 9(11), 5880–5890.
- Taniguchi, K., & Karin, M. (2018). NF- κ B, inflammation, immunity and cancer: Coming of age. *Nature Reviews Immunology*, 18(5), 309–324.
- Teng, H., Zheng, Y., Cao, H., Huang, Q., Xiao, J., & Chen, L. (2021). Enhancement of bioavailability and bioactivity of diet-derived flavonoids by application of nanotechnology: A review. *Critical Reviews in Food Science and Nutrition*, 1–16.
- Tomas, M., Beekwilder, J., Hall, R. D., Sagdic, O., Boyacioglu, D., & Capanoglu, E. (2017). Industrial processing versus home processing of tomato sauce: Effects on phenolics, flavonoids and in vitro bioaccessibility of antioxidants. *Food Chemistry*, 220, 51–58.
- Toro-Uribe, S., López-Giraldo, L. J., Alvarez-Rivera, G., Ibáñez, E., & Herrero, M. (2019). Insight of stability of procyanidins in free and liposomal form under an in vitro digestion model: Study of bioaccessibility, kinetic release profile, degradation, and antioxidant activity. *Journal of Agricultural and Food Chemistry*, 67(7), 1990–2003.
- Wan, Y., Wang, F., Yuan, J., Li, J., Jiang, D., Zhang, J., et al. (2019). Effects of dietary fat on gut microbiota and faecal metabolites, and their relationship with cardiometabolic risk factors: A 6-month randomised controlled-feeding trial. *Gut*, 68 (8), 1417–1429.
- Wang, X., Xia, K., Yang, X., & Tang, C. (2019). Growth strategy of microbes on mixed carbon sources. *Nature Communications*, 10(1), 1–7.
- Wang, F., Zhao, C., Tian, G., Wei, X., Ma, Z., Cui, J., et al. (2020). Naringin alleviates Atherosclerosis in ApoE $^{-/-}$ Mice by regulating cholesterol metabolism involved in gut microbiota remodeling. *Journal of Agricultural and Food Chemistry*, 68(45), 12651–12660.
- Wang, M., Zhao, H., Wen, X., Ho, C. T., & Li, S. (2021). Citrus flavonoids and the intestinal barrier: Interactions and effects. *Comprehensive Reviews in Food Science and Food Safety*, 20(1), 225–251.
- Williamson, G., Kay, C. D., & Crozier, A. (2018). The bioavailability, transport, and bioactivity of dietary flavonoids: A review from a historical perspective. *Comprehensive Reviews in Food Science and Food Safety*, 17(5), 1054–1112.
- Wu, H., & Ballantyne, C. M. (2020). Metabolic inflammation and insulin resistance in obesity. *Circulation Research*, 126(11), 1549–1564.
- Xiao, J. (2022). Recent advances in dietary flavonoids for management of type 2 diabetes. *Current Opinion in Food Science*, 44, Article 100806.
- Yan, R., & Zhang, X. (2022). Analysis of *Cyclocarya paliurus* flavonoids modulation on the physiology and gene expression in *Enterococcus faecalis* under bile salt stress. *Letters in Applied Microbiology*, 74(4), 555–563.
- Yan, X., Yang, C., Lin, G., Chen, Y., Miao, S., Liu, B., et al. (2019). Antidiabetic potential of green seaweed Enteromorpha prolifera flavonoids regulating insulin signaling pathway and gut microbiota in type 2 diabetic mice. *Journal of Food Science*, 84(1), 165–173.
- Yang, B., Liu, H., Yang, J., Gupta, V. K., & Jiang, Y. (2018). New insights on bioactivities and biosynthesis of flavonoid glycosides. *Trends in Food Science & Technology*, 79, 116–124.
- Yang, J., Wang, X., Zhang, G., Ma, L., Wei, T., Zhao, Y., et al. (2021). Comparative study of inhibition mechanisms of structurally different flavonoid compounds on α -glucosidase and synergistic effect with acarbose. *Food Chemistry*, 347, Article 129056.
- Yang, F., Zhu, W., Sun, S., Ai, Q., Edirisuriya, P., & Zhou, K. (2020). Isolation and structural characterization of specific bacterial β -glucuronidase inhibitors from Noni (*Morinda citrifolia*) fruits. *Journal of Natural Products*, 83(4), 825–833.
- T.S. Yeh C. Yuan A. Ascherio B.A. Rosner W.C. Willett D. Blacker Long-term dietary flavonoid intake and subjective cognitive decline in US men and women Neurology 97 10 2021 e1041.
- Yu, Y., Li, Z., Cao, G., Huang, S., & Yang, H. (2019). Bamboo leaf flavonoids extracts alleviate oxidative stress in HepG2 cells via naturally modulating reactive oxygen species production and Nrf2-mediated antioxidant defense responses. *Journal of Food Science*, 84(6), 1609–1620.
- Yu, Y., Pei, F., & Li, Z. (2022). Orientin and vitexin attenuate lipopolysaccharide-induced inflammatory responses in RAW264.7 cells: A molecular docking study, biochemical characterization, and mechanism analysis. *Food Science and Human Wellness*, 11(5), 1273–1281.
- Yuan, J., Che, S., Zhang, L., & Ruan, Z. (2021). Reparative effects of ethanol-induced intestinal barrier injury by flavonoid luteolin via MAPK/NF- κ B/MLCK and Nrf2 signaling pathways. *Journal of Agricultural and Food Chemistry*, 69(14), 4101–4110.
- Yuan, L., Li, X., He, S., Gao, C., Wang, C., & Shao, Y. (2018). Effects of natural flavonoid isoorientin on growth performance and gut microbiota of mice. *Journal of Agricultural and Food Chemistry*, 66(37), 9777–9784.
- Zeng, X., Yao, H., Zheng, Y., Chen, T., Peng, W., Wu, H., et al. (2019). Metabolite profiling of naringin in rat urine and feces using stable isotope-labeling-based liquid chromatography-mass spectrometry. *Journal of Agricultural and Food Chemistry*, 68 (1), 409–417.
- Zhang, F., Ma, Z., Qiao, Y., Wang, Z., Chen, W., Zheng, S., et al. (2022). Transcriptome sequencing and metabolomics analyses provide insights into the flavonoid biosynthesis in *Torreya grandis* kernels. *Food Chemistry*, 374, Article 131558.
- Zhang, B., Sun, W., Yu, N., Sun, J., Yu, X., Li, X., et al. (2018). Anti-diabetic effect of baicalein is associated with the modulation of gut microbiota in streptozotocin and high-fat-diet induced diabetic rats. *Journal of Functional Foods*, 46, 256–267.
- Zhang, M., Zhu, S., Yang, W., Huang, Q., & Ho, C. T. (2021). The biological fate and bioefficacy of citrus flavonoids: Bioavailability, biotransformation, and delivery systems. *Food & Function*, 12(8), 3307–3323.
- Zhao, S., Tian, G., Zhao, C., Li, C., Bao, Y., DiMarco-Crook, C., et al. (2018). The stability of three different citrus oil-in-water emulsions fabricated by spontaneous emulsification. *Food Chemistry*, 269, 577–587.
- Zhou, D., Bai, Z., Guo, T., Li, J., Li, Y., Hou, Y., et al. (2022). Dietary flavonoids and human top-ranked diseases: The perspective of in vivo bioactivity and bioavailability. *Trends in Food Science & Technology*, 120, 374–386.
- Zhu, Z., Han, Y., Ding, Y., Zhu, B., Song, S., & Xiao, H. (2021). Health effects of dietary sulfated polysaccharides from seafoods and their interaction with gut microbiota. *Comprehensive Reviews in Food Science and Food Safety*, 20(3), 2882–2913.
- Zmora, N., Suez, J., & Elinav, E. (2019). You are what you eat: Diet, health and the gut microbiota. *Nature Reviews Gastroenterology & Hepatology*, 16(1), 35–56.